

A Novel $SO(3)$ Picture for Quantum Searching

Gui Lu Long^{1,2,3}, Chang Cun Tu¹, Yan Song Li¹, Wei Lin Zhang¹ and Hai Yang Yan¹

¹*Department of Physics, Tsinghua University, Beijing 100084, China*

²*Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China*

³*Center of Nuclear Theory, National Laboratory of Heavy Ion Accelerator, Chinese Academy of Sciences, Lanzhou, 730000, China*

Abstract

An $SO(3)$ picture of the generalized Grover's quantum searching algorithm, with arbitrary unitary transformation and with arbitrary phase rotations, is constructed. In this picture, any quantum search operation is a rotation in a 3 dimensional space. Exact formulas for the rotation angle and rotational axis are given. The probability of finding the marked state is just $(z + 1)/2$, where z is the z -component of the state vector. Exact formulas for this probability is easily obtained. The phase matching requirement and the failure of algorithm when phase mismatches are clearly explained.

03.67-a, 03.67.Lx, Quantum searching, Phase matching, $SO(3)$ group

Grover's quantum search algorithm [1,2] is one of the most celebrated quantum computing algorithms. It has been shown that the algorithm is optimal [3]. The algorithm can be generalized to arbitrary initial amplitude distribution [4]. It has many important applications, for instance, in the Simon problem [5] and quantum counting [6]. In the case where multiple marked state is involved, it can even search the data by just one query [7]. Recently, it has been generalized to an arbitrarily entangled initial states [8]. Since Grover's algorithm involves only simple gate operations, it has been realized in 2 qubits [9–11], and 3 qubit NMR systems [12].

Grover's original algorithm has a simple geometric interpretation [2,13,15]. When the Hadmard transformation is substituted by any arbitrary unitary transformation, it has been shown there is an $SU(2)$ group structure in the generalized algorithm [2,15]. However, when generalizing the algorithm to arbitrary phase rotations, phase matching is vital [17,18]. In [18] we have given an approximate formula for the amplitude of the marked state. But it is difficult to understand the phase matching requirement, as it is contrary to what one expects from an continuity argument.

In this Letter, we give a novel $SO(3)$ picture of the generalized quantum search algorithm by exploiting the relation between $SO(3)$ and $SU(2)$. In this $SO(3)$ picture the process of quantum search is crystalline transparent. The behavior of the algorithm with phase matching or mismatching are clearly understood. This helps us to understand the various aspects of the algorithm, and to further develop the algorithm.

The operator for quantum search [2] can be written as $Q = -I_\gamma U^{-1} I_\tau U$, where $|\tau\rangle$ is the marked state, $|\gamma\rangle$ is the prepared state, usually $|\gamma\rangle = |0\rangle$. For arbitrary phase rotations, $I_\gamma = I - (-e^{i\theta} + 1) |\gamma\rangle\langle\gamma|$, $I_\tau = I - (-e^{i\phi} + 1) |\tau\rangle\langle\tau|$. In the basis where $|1\rangle = U^{-1}|\tau\rangle$, $|2\rangle = -(|\gamma\rangle - U_{\tau\gamma}U^{-1}|\tau\rangle)/\sqrt{1 - |U_{\tau\gamma}|^2}$, Q can be written as

$$Q = \begin{pmatrix} -e^{-i\frac{\phi}{2}}(\cos\frac{\theta}{2} + i\cos 2\beta\sin\frac{\theta}{2}) & -ie^{-i\frac{\phi}{2}}\sin 2\beta\sin\frac{\theta}{2} \\ -ie^{i\frac{\phi}{2}}\sin 2\beta\sin\frac{\theta}{2} & -e^{i\frac{\phi}{2}}(\cos\frac{\theta}{2} - i\cos 2\beta\sin\frac{\theta}{2}) \end{pmatrix}, \quad (1)$$

where we have written $U_{\tau\gamma} = e^{i\xi}\sin\beta$ (in Grover's original algorithm, $U_{\tau\gamma} = \frac{1}{\sqrt{N}}$, $\xi = 0$, $\sin\beta = \frac{1}{\sqrt{N}}$), and an overall phase factor has been neglected.

It is easy to check that $\det(Q) = 1$, and Q is an element of the $SO(3)$ group. As is well known, each unitary matrix u in $SU(2)$ group corresponds to a rotation R_u in $SO(3)$ group [16]. Here operator Q corresponds to the rotation,

$$\begin{pmatrix} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{pmatrix} \quad (2)$$

where $R_{11} = \cos\phi(\cos^2 2\beta\cos\theta + \sin^2 2\beta) + \cos 2\beta\sin\theta\sin\phi$, $R_{12} = \cos 2\beta\cos\phi\sin\theta - \cos\theta\sin\phi$, $R_{13} = -\cos\phi\sin 4\beta\sin^2\frac{\theta}{2} + \sin 2\beta\sin\theta\sin\phi$, $R_{21} = -\cos(2\beta)\cos\phi\sin\theta + (\cos^2\frac{\theta}{2} - \cos 4\beta\sin^2\frac{\theta}{2})\sin\phi$, $R_{22} = \cos\theta\cos\phi + \cos 2\beta\sin\theta\sin\phi$, $R_{23} = -\cos\phi\sin 2\beta\sin\theta - \sin 4\beta\sin^2\frac{\theta}{2}\sin\phi$, $R_{31} = -\sin 4\beta\sin^2\frac{\theta}{2}$, $R_{32} = \sin 2\beta\sin\theta$, $R_{33} = \cos^2 2\beta + \cos\theta\sin^2 2\beta$.

A spinor in $SU(2)$ which describes the state of the quantum computer, $\Psi = \begin{pmatrix} a + bi \\ c + di \end{pmatrix}$ corresponds to a vector in R^3

$$\mathbf{r} = \Psi^\dagger \sigma \Psi = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2(ac + bd) \\ 2(-bc + ad) \\ a^2 + b^2 - c^2 - d^2 \end{pmatrix}. \quad (3)$$

The probability of finding the marked state is $P = a^2 + b^2 = (z + 1)/2$. The z component of the polarization vector is a measure of the probability. For instance, the evenly distribution state $\Psi_o = (\frac{1}{\sqrt{N}}, \frac{\sqrt{N-1}}{\sqrt{N}})^\dagger$, corresponds to vector $\mathbf{r}_o = (2\sqrt{1 - \frac{1}{N}}\sqrt{\frac{1}{N}}, 0, -1 + \frac{2}{N})^T$, which is nearly parallel to the $-z$ axis when N is large. The marked state $\psi_a = (1, 0)^\dagger$, corresponds to $\mathbf{r}_a = (0, 0, 1)^T$, which is on the $+z$ axis. Thus the process of quantum searching in the $SO(3)$ picture is to rotate the state vector from a position nearly parallel to $-z$ axis to $+z$ axis.

The rotational axis of (3) can be found by solving the eigen-value problem, $R_u \mathbf{l} = \mathbf{l}$. This gives $\mathbf{l} = \left(\cot \frac{\phi}{2} \quad 1 \quad -\cot 2\beta \cot \frac{\phi}{2} + \cot \frac{\theta}{2} \csc 2\beta \right)^T$. Each iteration of Q rotates about this axis an angle

$$\alpha = \arccos \left[\frac{1}{4} (\cos 4\beta + 3) \cos \theta \cos \phi + \sin^2 2\beta \left(\frac{1}{2} \cos \phi - \sin^2 \frac{\theta}{2} \right) + \cos 2\beta \sin \theta \sin \phi \right], \quad (4)$$

about the rotational axis. In Grover's original algorithm, $\theta = \phi = \pi$, the rotation axis is exactly the y -axis, and the rotational angle is equal to the maximum value of 4β (remember the relation between $SU(2)$ and $SO(3)$, this corresponds an angle of 2β in the $SU(2)$). The state vector $\mathbf{r}$ is being rotated within the $x - z$ plane from approximately $-z$ to $+z$ axis, where the marked state achieves maximum probability amplitude. The number of step requires to reach $+z$ axis is $\frac{\pi - 2\beta}{\alpha} \approx 0.785\sqrt{N} - 0.5 \approx 0.785\sqrt{N}$. The trace of tip of the state vector is shown in Fig.1.

In the most general case with arbitrary θ and ϕ , the trace of the tip of state vector $\mathbf{r}$ is a circle. The state vector spans a cone with the top at the origin. During the rotation, the vector $\mathbf{r} - \mathbf{r}_o$ is orthogonal to the rotational axis $\mathbf{l}$ at any time: $(\mathbf{r} - \mathbf{r}_o) \cdot \mathbf{l} = 0$. If the state vector passes through $+z$ axis, that is $\mathbf{r} = (0, 0, 1)^T$ be in the trace, by solving equation $\mathbf{r} - \mathbf{r}_a \cdot \mathbf{l} = 0$, we have $\cot \frac{\phi}{2} = \cot \frac{\theta}{2}$, or $\phi = \theta$, the phase matching requirement which has been found in an approximate manner. However, the rotational axis is now $\mathbf{l} = \left(\cos \frac{\phi}{2} \quad \sin \frac{\phi}{2} \quad \cos \frac{\phi}{2} \tan \beta \right)^T$, which is no longer the y axis. The rotation angle is

$$\alpha = \arccos \{ 2[(\cos 2\beta - 1) \sin^2 \frac{\phi}{2} + 1]^2 - 1 \}. \quad (5)$$

If N is very large, $\mathbf{l} \approx \left(\cos \frac{\phi}{2} \quad \sin \frac{\phi}{2} \quad 0 \right)$, which is in the $x - y$ plane, and the initial state vector is nearly the $-z$ axis. The trace of tip of the state vector is a circle in the $x - z$ plane. Each iteration rotates the state vector an angle α given by (11). To first order in β , $\alpha \approx 4\beta \sin \frac{\phi}{2}$, which corresponds a rotation of $2\beta \sin \frac{\phi}{2}$ in $SU(2)$. The number of steps requires to search the marked state is larger than that in the original version, as given in [18]. However in this case, the centre of the circle is no longer the origin. The state vector can pass the $+z$ axis, that is, it can reach the marked state with near certainty, but not the $-z$ where the amplitude of the marked state is zero. This has clearly been demonstrated in the numerical calculation in Ref. [18].

When $\theta \neq \phi$, the trace the tip of the state vector is still a circle. But it is very tilted. In Figure 2, it is drawn for the case of $\theta = \frac{\pi}{2}$, $\phi = \frac{\pi}{10}$. Here we see the rotating axis is nearly the z axis, the circle span by the state vector tip is nearly parallel to the $x - y$ plane. Therefore the amplitude of the marked state can not reach 1, neither can it reach zero. This explains naturally the intriguing narrowly bounded behavior of the algorithm we have found in Ref. [17].

To summarize, we have given a novel $SO(3)$ interpretation of the quantum search algorithm. In this picture, the effect of quantum search is clearly displayed. In particular, the phase-mismatching are clearly understood. This throws new light on the algorithm, and we hope it helpful for further development of the algorithm.

REFERENCES

- [1] Lov K. Grover, Phys. Rev. Lett. **79**(1997) 325.
- [2] L. K. Grover, Phys. Rev. Lett. **79** (1997) 4709.
- [3] C. Zalka, Lanl-eprint quant-ph/9711070.
- [4] M. Boyer, G. Brassard, P. Hoyer, A. Tapp, Fortsch. Phys. **46** (1998) 493.
- [5] G/ Brassard and P. Hoyer, quant-ph/9704027.
- [6] G. Brassard, P. Hoyer and A. Tapp, quant-ph/9805082.
- [7] D. P. Chi and J. Kim, quant-ph/9708005.
- [8] A. Carlilni and A. Hosoya, quant-ph/9909089.
- [9] I. L. Chuang, N. Gershenfeld and M. Kubinee, Phys. Rev. Lett. **80** (1998) 3408.
- [10] J. A. Jones, M. Mosca and R. H. Hansen, Nature, **393** (1998) 344.
- [11] Fiping Fu, Li Xiao, Jum Luo and Xizhi Zheng, J. Chin. Mag. Res., **16** (1999) 341.
- [12] L M. K. Vandersypen et. al. Lanl-eprint quant-ph/9910075.
- [13] R. Jozsa, quant-ph/9901021.
- [14] A. Kumar Pati, quant-ph/9807067.
- [15] Sixia Yu, Chang-Pu Sun, quant-ph/990375.
- [16] Q. Z. Han and H. Z. Sun, Group Theory, Peking University Press, 1987; H. Z. Sun and Q. Z. Han, Lie Algebras and Lie Superalgebras and Their Applications in Physics, Peking University Press, 1999.
- [17] G. L. Long et. al., Commun. Theor. Phys. **32** (1999) 335, also in quant-ph/9904077.
- [18] G. L. Long et. al., Phys. Lett. A **262** (1999) 27, also in quant-ph/996020.
- [19] C. Zalka, quant-ph/9902049.

FIGURES




FIG. 1. The trace of the vector state tip when phase matching is satisfied.




FIG. 2. 3D plot of the trace of the vector state tip when phase mismatches.